illustra™ ExoProStar™

Enzymatic PCR and Sequencing Clean-Up

Codes: US78210

US78211 US78212 US78220 US78225

Description

The ExoProStar PCR and Sequence Reaction Clean-Up Kit is designed for direct clean-up of unincorporated primers and dNTPs from PCR reactions prior to DNA sequencing. illustra ExoProStar improves digestion efficiency without degradation of the target PCR product which remains completely intact and ready for immediate use in manual or automated processes.

Legal

GE, imagination at work and GE Monogram are trademarks of General Electric Company. illustra and ExoProStar are trademarks of General Electric Company or one of its subsidiaries. Exonuclease I/Alkaline Phosphatase method of use is covered by one or more of the following US patent numbers: 5,756,285 and 5,741,676 in the name of GE Healthcare Bio-Sciences Corp.

All goods and services are sold subject to the terms and conditions of sale of the company within GE Healthcare which supplies them. A copy of these terms and conditions is available on request. Contact your local GE Healthcare representative for the most current information.

 $\hbox{@ 2013 General Electric Company - All rights reserved}$

First published Mar 2013

For your local office contact information, visit http://www.gelifesciences.com/contact

GE Healthcare UK Limited Amersham Place Little Chalfont, Buckinghamshire, HP7 9NA, UK

http://www.gelifesciences.com



Recommended Protocol

- **1.** Remove the Alkaline Phosphatase and Exonuclease I from the freezer and keep on ice whilst preparing the reaction.
- 2. Take a 5 µl aliquot of the completed PCR reaction.
- 3. Add $1 \mu l$ of Alkaline Phosphatase to the reaction mix.
- **4.** Add 1 μ l of Exonuclease I to the reaction mix.
- 5. Incubate at 37°C for 15 minutes.
- **6.** Incubate at 80°C for 15 minutes to inactivate the enzymes.
- **7.** The PCR product is now ready for use in sequencing or other down-stream applications as required.

This reaction is scalable for larger quantities of PCR reaction mix. Simply add Alkaline Phosphatase and Exonuclease I in proportion to the volume of PCR reaction mix required.

